

JONATHAN E. FIELDING, M.D., M.P.H.
Director and Health Officer

LAURENE MASCOLA, M.D., M.P.H.
Chief, Acute Communicable Disease Control

Immunization Program

MICHELLE T. PARRA, Ph.D. Director
A. NELSON EL AMIN, M.D., M.P.H. Medical Director

3530 Wilshire Boulevard, Suite 700
Los Angeles, California 90010
TEL (213) 351-7800 • FAX (213) 351-2780

www.lapublichealth.org

April 10, 2008

BOARD OF SUPERVISORS

Gloria Molina
First District

Yvonne B. Burke
Second District

Zev Yaroslavsky
Third District

Don Knabe
Fourth District

Michael D. Antonovich
Fifth District

MEASLES ALERT FOR LOS ANGELES COUNTY

There has been one case of measles identified in the West Valley area of Los Angeles County. Although this is the first reported measles case in Los Angeles since 2006, there has recently been a cluster of measles cases reported in San Diego and in our neighboring State of Arizona. Measles is a highly communicable disease that can have serious complications. Infected persons are capable of transmitting measles for at least four days before the beginning of the rash to four days after its appearance.

- **Consider measles when evaluating any patient who has an acute rash illness with fever.** Symptoms usually begin 10-12 days (up to 18 days) after exposure with a prodrome of fever as high as 105 degrees, malaise, cough, runny nose, and conjunctivitis. Two to four days following the onset of the prodrome, a maculopapular rash develops. The rash usually begins around the ears and the hairline and then spreads down to cover the face, trunk, arms, and legs.
- **IMMEDIATELY report any suspect cases in Los Angeles County to the Morbidity Central Reporting Unit at 888-397-3993 (phone) or 888-397-3778 (fax).** Do NOT wait for laboratory confirmation before reporting a suspect case. During non-business hours (before 7:30 am, after 5:00 pm, or weekends) call 213-974-1234. For cases among residents of the Cities of Long Beach or Pasadena, contact their respective health departments.
- **Obtain blood specimens for serologic confirmation of the diagnosis.** Blood specimens should be obtained at the time of clinical presentation and be tested for both measles IgM and IgG antibodies. Additionally, specimens for measles virus culture should be collected and transported in a tube of viral transport media. If within 4 days of rash onset, collect both a nasopharyngeal swab and a urine specimen. If within 5-7 days of rash onset, collect urine specimen only.
- **Respiratory and airborne precautions should be taken to prevent transmission.** Suspect measles cases needing medical attention should not be allowed in patient waiting areas. They should be masked and placed immediately in an examination room, with the door closed. The examination room in which a patient with suspected measles or confirmed measles was examined should not be used again for two hours. With hospitalized patients, respiratory isolation is recommended for 4 days after the onset of rash. Listings of all exposed patients and staff will be required by the health department from any medical facility with a case of measles.
- **Recommend MMR vaccination as post-exposure prophylaxis for susceptible contacts.** MMR vaccine given within 72 hours of exposure can prevent measles. For susceptible persons who cannot be vaccinated, immune globulin (IG) can be given within 6 days of exposure to prevent measles or modify the disease if it occurs.

For technical assistance or additional information about measles, please call the Los Angeles County Immunization Program's Surveillance Unit at (213) 351-7800.